

NON VANISHING OF CENTRAL VALUES OF MODULAR L-FUNCTIONS FOR HECKE EIGENFORMS OF LEVEL ONE

D. CHOI AND Y. CHOIE

ABSTRACT. Let $F(z) = \sum_{n=1}^{\infty} a(n)q^n$ be a newform of weight $2k$ and level N with a trivial character, and assume that $F(z)$ is a non-zero eigenform of all Hecke operators. For $x > 0$, let

$$\mathcal{N}_F(x) := |\{D \text{ fundamental} \mid |D| < x, (D, N) = 1, L(F, D, k) \neq 0\}|.$$

A based on the Goldfeld's conjecture one expects to have $\mathcal{N}_F(x) \gg x$ ($x \rightarrow \infty$). Kohnen [10] showed that if $k \geq 6$ is a even integer, then for $x \gg 0$ there is a normalized Hecke eigenform F of level 1 and weight $2k$ with the property that

$$\mathcal{N}_F(x) \gg_k x \quad (x \rightarrow \infty).$$

In this paper, we extend the result in [10] to the case when k is any integer, in particular when k is odd. So, we obtain that, when the level is 1, for each integer $2k$ such that the dimension of cusp forms of weight $2k$ is not zero, there is a normalized Hecke eigenform F of weight $2k$ satisfying $\mathcal{N}_F(x) \gg x$ ($x \rightarrow \infty$).

January 28, 2010

1. Introduction and statement of result

Let $F(z) = \sum_{n=1}^{\infty} a(n)q^n \in S_{2k}(N, \chi_0)$ be a newform of weight $2k$ and level N with a trivial character χ_0 , and assume that $F(z)$ is a non-zero eigenform of all Hecke operators. For a fundamental discriminant D , that is $D = 1$ or the discriminant of a quadratic field, we define the L -function $L(F, D, s)$ of F twisted with the quadratic character $(\frac{D}{\bullet})$ by

$$\sum_{n \geq 1} \left(\frac{D}{n} \right) \frac{a(n)}{n^s}.$$

In this paper, we consider the central values of L -functions $L(F, D, k)$.

It is well-known by Waldspurger [18] that the central critical values $L(F, D, k)$ are essentially proportional to the squares of Fourier coefficients of the modular form of weight $k + \frac{1}{2}$ corresponding to F under Shimura correspondence.

On the other hand, for $x > 0$, consider the set

$$\mathcal{N}_F(x) := |\{D \text{ fundamental} \mid |D| < x, (D, N) = 1, L(F, D, k) \neq 0\}|.$$

2000 *Mathematics Subject Classification.* 11F67, 11F37.

Key words and phrases. modular L-functions, quadratic twist, central L-values.

As based on the Goldfeld's conjecture one expected to have

$$(1.1) \quad \mathcal{N}_F(x) \gg x \quad (x \rightarrow \infty)$$

(In [5] Goldfeld conjectured that

$$\begin{aligned} & \sum_{D \text{ fundamental, } |D| < x, \gcd(D, N)=1} \text{ord}_{s=k} L(F, D, s) \\ & \approx \frac{1}{2} |\{D \text{ fundamental} \mid |D| < x, \gcd(D, N) = 1\}| \quad (x \rightarrow \infty). \end{aligned}$$

In [14] using the theory of Galois representations of modular forms together with the results of Friedberg and Hoffstein [4], the following was proved

$$\mathcal{N}_F(x) \gg_F \frac{x}{\log x} \quad (x \rightarrow \infty).$$

In [7] K. James gave an example of a form F , by looking at the difference of two special ternary theta series of level 56, satisfying $\mathcal{N}_F(x) \gg_F x \quad (x \rightarrow \infty)$. Furthermore, Kohnen [10] showed that if $k \geq 6$ is an even integer, then there is a normalized Hecke eigenform F of level 1 and weight $2k$ with the property that $\mathcal{N}_F(x) \gg_k x$ for $x \gg 0$; in particular, it was shown that $\mathcal{N}_\Delta(x) \gg x \quad (x \rightarrow \infty)$ where Δ is Ramanujan's Δ -function of weight 12. More precisely, Kohnen proved that if $N_{k, \Gamma_1}^+(x)$ is the number of fundamental discriminant D , $0 < D < x$, such that there exists a normalized Hecke eigenform $F \in S_{2k}$ satisfying $L(F, D, k) \neq 0$, then for any positive ϵ

$$\mathcal{N}_{k, \Gamma_1}^+(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

Here, S_{2k} is the space of cusp forms of weight $2k$ on $\Gamma_1 = SL(2, \mathbb{Z})$. In this paper, we extend the result [10] to the case when k is any integer, in particular when k is odd. To obtain these results, we refine the argument of Kohnen by using an isomorphism [8] from the spaces of modular forms of integral weight to the Kohnen plus space of half integral weight modular forms.

Let $N_{k, \Gamma_1}(x)$ denote the number of fundamental discriminants D , $|D| < x$, such that there exists a normalized Hecke eigenform $F \in S_{2k}$ satisfying $L(F, D, k) \neq 0$, then we have the following:

Theorem 1.1. *Suppose that $k \geq 9$ is odd. Then for any positive ϵ*

$$\mathcal{N}_{k, \Gamma_1}(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

Let $d_k := \dim(S_{2k})$. With the result of Kohnen [10] in which covers the case of $k \equiv 0 \pmod{2}$, we state the following result:

Theorem 1.2. *Suppose that k is a positive integer such that $d_k \geq 1$. Then for any $\epsilon > 0$*

$$\mathcal{N}_{k,\Gamma_1}(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

Theorem 1.2 immediately implies that for each integer k such that $d_k > 0$ there is a Hecke eigenform $F \in S_{2k}$ satisfying (1.1).

Corollary 1.3. *Suppose that k is an integer such that $d_k \geq 1$. Then there exists a Hecke eigenform $F(z) \in S_{2k}$ such that for any positive ϵ*

$$\mathcal{N}_F(x) \geq \frac{1}{d_k} \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

Recently, Farmer and James [3] proved that the characteristic polynomial of the Hecke operator T_2 on S_{2k} is irreducible over $\mathbb{Q}$ for $k \leq 1,000$. If K is the field obtained from $\mathbb{Q}$ by adjoining the eigenvalues of T_2 , then the Galois group $G = \text{Gal}(K/\mathbb{Q})$ operates transitively on the set of normalized Hecke eigenforms in S_{2k} . Using Theorem 1.1 and the known fact [15] that $L(F^\sigma, D, k)_{\text{alg}} = L(F, D, k)_{\text{alg}}^\sigma$ for all $\sigma \in G$, we have that every Hecke eigenform $F \in S_{2k}$ satisfies (1.1) for each integers k , $6 \leq k \leq 1,000$. Here, “alg” means “algebraic part”.

Theorem 1.4. *Suppose that $6 \leq k \leq 1,000$ is an integer. Then every normalized Hecke eigenform F in S_{2k} satisfies*

$$(1.2) \quad \mathcal{N}_F(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

Remark 1.5. Maeda ([6] Conjecture 1.2) made a conjecture that the Hecke algebra of S_{2k} over $\mathbb{Q}$ is simple, and that its Galois closure over $\mathbb{Q}$ has Galois group G the full symmetric group. The conjecture implies that there is a single Galois orbit of Hecke eigenforms in S_{2k} . Thus, Maeda’s conjecture implies that every normalized Hecke eigenform F in S_{2k} satisfies (1.2).

2. Preliminaries

Let $q := e^{2\pi iz}$, where z is in the complex upper half plane $\mathbb{H}$. For an integer $k \geq 2$ recall the normalized Eisenstein series $E_{2k}(z) := 1 - \frac{4k}{B_{2k}} \sum_{n \geq 1} \sigma_{k-1}(n) q^n$ of weight $2k$, and for an integer $r \geq 1$ let $H_{r+\frac{1}{2}}(z) = \sum_{N=0}^{\infty} H(r, N) q^N$ be the Cohen-Eisenstein series of weight $r + \frac{1}{2}$ on $\Gamma_0(4)$ (see [1]). Here, for each positive integer N , define

$$h(r, N) = \begin{cases} (-1)^{\lfloor \frac{r}{2} \rfloor} (r-1)! N^{r-\frac{1}{2}} 2^{1-r} \pi^{-r} L(r, \chi_{(-1)^r N}) & \text{if } (-1)^r N \equiv 0 \text{ or } 1 \pmod{4}, \\ 0 & \text{if } (-1)^r N \equiv 2 \text{ or } 3 \pmod{4}, \end{cases}$$

where χ_D denotes the character $\chi_D(d) = (\frac{D}{d})$. Furthermore, for $N \geq 1$, define

$$H(r, N) = \begin{cases} \sum_{d^2|N} h(r, \frac{N}{d^2}) & \text{if } (-1)^r N \equiv 0 \text{ or } 1 \pmod{4}, \\ \zeta(1-2r) & \text{if } N = 0, \\ 0 & \text{otherwise.} \end{cases}$$

The followings are proved in [1]:

Proposition 2.1. (1) For $r \geq 2$, Cohen-Eisenstein series $H_{r+\frac{1}{2}}(z)$ is a modular form of weight $r + \frac{1}{2}$ on $\Gamma_0(4)$ and it is in Kohnen plus condition, that is,

$$H(r, N) = 0 \text{ if } (-1)^r N \not\equiv 0, 1 \pmod{4}.$$

(2) Let a and b be integers with $a \geq 1$. Suppose that $-b$ is a quadratic non residue of a . Then the function

$$G_{a,b}(z) := \sum_{N \equiv b \pmod{a}} H(1, n) q^n$$

is a modular form of weight $\frac{3}{2}$ and character a over $\Gamma_0(A)$, where we can take $A = 4a^2$, and furthermore $A = a^2$ if a is even.

For a nonnegative integer k denote $M_{k+\frac{1}{2}}(\Gamma_0(4))$ as the usual complex vector space of cusp forms of weight $k + \frac{1}{2}$ on $\Gamma_0(4)$ with the trivial character. In [8], Kohnen introduces the plus space $M_{k+\frac{1}{2}}^+(\Gamma_0(4))$ of modular forms $g(z)$ of weight $k + \frac{1}{2}$ on $\Gamma_0(4)$ with a Fourier expansion of the form

$$g(z) = \sum_{(-1)^k n \equiv 0, 1 \pmod{4}} c(n) q^n.$$

and proved the following isomorphism (Proposition 1 in [8]).

Proposition 2.2. Let M_k be the space of modular forms of weight k on Γ_1 . If k is even, then the spaces $M_k \oplus M_{k-2}$ and $M_{k+\frac{1}{2}}^+(\Gamma_0(4))$ are isomorphic under the map

$$(f(z), h(z)) \mapsto f(4z)\theta(z) + h(4z)H_{\frac{5}{2}}(z),$$

where $\theta(z) = 1 + 2 \sum_{n=1}^{\infty} q^{n^2}$. If k is odd, then the spaces $M_{k-3} \oplus M_{k-5}$ and $M_{k+\frac{1}{2}}^+(\Gamma_0(4))$ are isomorphic under the map

$$(f(z), h(z)) \mapsto f(4z)H_{\frac{7}{2}}(\tau) + h(4z)H_{\frac{11}{2}}(z).$$

For $k \geq 2$ we have $M_{k+\frac{1}{2}}^+(\Gamma_0(4)) = \mathbb{C}H_{k+\frac{1}{2}} \oplus S_{k+\frac{1}{2}}^+(\Gamma_0(4))$.

The results of [11], [9], and [18] connect the coefficients of Hecke eigenforms of half-integral weight to the central L -values of twists of integral weight Hecke eigenforms. More precisely, suppose that $f(z) = \sum_{n=1}^{\infty} a(n)q^n \in S_{2k}$ is a normalized Hecke eigenform and that $g(z) = \sum_{n=1}^{\infty} c(n)q^n \in S_{k+\frac{1}{2}}^+(\Gamma_0(4))$ is a Hecke eigenform with the same Hecke eigenvalues as those of f . Here,

$$S_{k+\frac{1}{2}}^+(\Gamma_0(4)) := \{g \in S_{k+\frac{1}{2}}(\Gamma_0(4)) \mid g(z) = \sum_{n \geq 1, (-1)^k n \equiv 0,1 \pmod{4}} c(n)q^n\}$$

Theorem 1 of [11] states the following.

Theorem 2.3. *Suppose that f and g are as above, D is a fundamental discriminant with $(-1)^k D > 0$, and $L(f, D, s)$ is the twisted L -series*

$$L(f, D, s) = \sum_{n=1}^{\infty} \left(\frac{D}{n}\right) a(n) n^{-s}.$$

Then

$$\frac{c(|D|)^2}{\langle g, g \rangle} = \frac{(k-1)!}{\pi^k} |D|^{k-\frac{1}{2}} \frac{L(f, D, k)}{\langle f, f \rangle}.$$

Here, $\langle g, g \rangle$ and $\langle f, f \rangle$ are the normalized Petersson scalar products

$$\begin{aligned} \langle g, g \rangle &= \frac{1}{6} \int_{\Gamma_0(4) \backslash \mathbb{H}} |g(z)|^2 y^{k-3/2} dx dy \\ \langle f, f \rangle &= \int_{\Gamma_1 \backslash \mathbb{H}} |f(z)|^2 y^{2k-2} dx dy. \end{aligned}$$

3. Proof of Theorem 1.1

For any function $f(z)$ on $\mathbb{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ and any positive integer d we define the operator U_d

$$(3.1) \quad (f|U_d)(z) := \frac{1}{d} \sum_{j=0}^{d-1} f\left(\frac{z+j}{d}\right).$$

Suppose that f has a Fourier expansion $f(z) = \sum a(n)q^n$. Then $(f|U_d)(z) = \sum a(nd)q^n$. If $g(z) = \sum_{n=0}^{\infty} c(n)q^n$ is in $M_{k+\frac{1}{2}}(\Gamma_0(4N))$, then the Hecke operator with the trivial character on $M_{k+\frac{1}{2}}(\Gamma_0(4N))$ is defined for odd primes ℓ by

$$(g|T(\ell^2, k))(z) := \sum_{n=0}^{\infty} \left(c(\ell^2 n) + \ell^{k-1} \left(\frac{(-1)^k n}{\ell} \right) c(n) + \left(\frac{(-1)^k}{\ell^2} \right) \ell^{2k-1} c\left(\frac{n}{\ell^2}\right) \right) q^n,$$

where $\left(\frac{\cdot}{\ell}\right)$ and $\left(\frac{\cdot}{\ell^2}\right)$ are Jacobi symbols, and $c\left(\frac{n}{\ell^2}\right) := 0$ if $\ell^2 \nmid n$. If g has integral coefficients, then one also has that

$$(3.2) \quad g|U_\ell \equiv g^\ell | T\left(\ell^2, \ell k + \frac{(\ell-1)}{2}\right) \pmod{\ell}.$$

For any positive integer d we define the operator V_d

$$(3.3) \quad (g|V_d)(z) := \sum_{n=0}^{\infty} c(n)q^{dn}.$$

Note that if ℓ is a prime, then

$$(3.4) \quad (g|V_\ell)(z) \equiv g(z)^\ell \pmod{\ell}.$$

For a Dirichlet character χ let

$$g \otimes \chi := \sum_{n=0}^{\infty} \chi(n) c(n) q^n.$$

The following proposition immediately implies our main theorems.

Proposition 3.1. *Suppose that $k \geq 8$ is an integer.*

(1) *If k is odd, then for any positive ϵ*

$$\mathcal{N}_{k,\Gamma_1}(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

(2) *If k is an even integer such that $d_k > 1$ or $k = 10$, then for any positive ϵ*

$$\mathcal{N}_{k,\Gamma_1}(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_\epsilon 0).$$

Proof For a positive even positive integer t , let

$$m(t) := \frac{1}{2} \left(t - 4 \left\lfloor \frac{t}{4} \right\rfloor \right).$$

For a non negative integer t we define a modular form $R_t(z)$:

$$R_t(z) := \begin{cases} E_4(4z)^{\lfloor \frac{t}{4} \rfloor - m(t)} E_6(4z)^{m(t)} & \text{if } t > 0, \\ 1 & \text{if } t = 0. \end{cases}$$

For any even positive integer t , we have

$$(3.5) \quad R_t(z) \equiv 1 \pmod{3}.$$

This is from the fact that if $k \geq 4$ is an even integer, then $E_{p-1}(z) \equiv 1 \pmod{p}$ for any prime p such that $k \equiv 0 \pmod{p-1}$ (see [12]).

First, we assume that k is odd. For each odd integers $k \geq 9$ define

$$(3.6) \quad \Phi_{k+\frac{1}{2}}(z) = 28H_{3+\frac{1}{2}}(z)R_{k-3}(z) - \frac{44}{3}H_{5+\frac{1}{2}}(z)R_{k-5}(z) := \sum_{n=1}^{\infty} \beta_k(n)q^n.$$

Then $\Phi_{k+\frac{1}{2}}(z)$ is in $S_{k+\frac{1}{2}}^+(\Gamma_0(4))$ by Proposition 2.2, and the Fourier coefficients of $\Phi_{k+\frac{1}{2}}(z)$ are 3-integral. On the other hand, for every odd $k \geq 9$, we have

$$\Phi_{k+\frac{1}{2}}(z) \equiv \Phi_{9+\frac{1}{2}}(z) (= 28H_{3+\frac{1}{2}}(z)R_6(z) - \frac{44}{3}H_{5+\frac{1}{2}}(z)R_4(z)) \pmod{3}.$$

Let

$$\begin{aligned} F(z) &:= (\Phi_{9+\frac{1}{2}}(z) - (\Phi_{9+\frac{1}{2}}(z)|U_3|V_3)) + (\Phi_{9+\frac{1}{2}}(z) - (\Phi_{9+\frac{1}{2}}(z)|U_3|V_3)) \otimes \left(\frac{\cdot}{3}\right) \\ &= \sum_{n \equiv 1 \pmod{3}} \beta_9(n) q^n. \end{aligned}$$

Recall that

$$G_{3,1}(z) = \sum_{n \equiv 1 \pmod{3}} H(1, n) q^n.$$

Proposition 2.1 implies that $G_{3,1}(z) = \sum_{n \equiv 1 \pmod{3}} H(1, n) q^n$ is a modular form of weight $\frac{3}{2}$ on $\Gamma_0(36)$ such that its coefficients are 3-integral.

Thus, by computing a few coefficient modulo 3 and using Sturm's bound in ([16]) we have

$$\begin{aligned} (3.7) \quad F(z) &\equiv \sum_{n \equiv 1 \pmod{3}} \beta_9(n) q^n \\ &\equiv 2q^4 + q^7 + q^{19} + 2q^{28} + 2q^{40} + q^{43} + 2q^{52} + q^{55} + 2q^{64} + q^{67} + q^{76} + \dots \\ &\equiv \sum_{n \equiv 1 \pmod{3}} H(1, n) q^n \equiv G_{3,1}(z) \pmod{3}. \end{aligned}$$

On the other hand, let $h(D) = H(1, D)$ be the class number of $\mathbb{Q}(\sqrt{D})$. It is known that for $D < 0$, $h(D) = -B_{1,(\frac{D}{\cdot})}$ (apart from $D = -3$ and -4) (for example, see [17]). Thus, we have

$$\beta_k(D) \equiv B_{1,(\frac{-D}{\cdot})} \equiv h(-D) \pmod{3},$$

for a fundamental discriminant $D > 1$ such that $D \equiv 1 \pmod{3}$.

Now let m and N be positive integers satisfying the condition:

(**) *If an odd prime p is a common divisor of m and N , then $p \mid N$ and $p^2 \nmid m$. Further if N is even, then (i) $4 \mid N$ and $m \equiv 1 \pmod{4}$ or (ii) $16 \mid N$ and $m \equiv 8$ or $12 \pmod{16}$.*

We denote by $N_2^-(x, m, N)$ the number of fundamental discriminants D with $-x < D < 0$ and $D \equiv m \pmod{N}$. The results of Davenport-Heilbronn [2] and Nakagawa-Horie [13] imply that for any positive number ϵ

$$(3.8) \quad \begin{aligned} &|\{\text{fundamental discriminants } D \equiv 1 \pmod{3} \mid 0 < D < x \text{ and } 3 \nmid h(-D)\}| \\ &\gg \left(\frac{1}{2} - \epsilon\right) (N_2^-(x, 1, 3)). \end{aligned}$$

Since $N_2^-(x, 1, 3) \sim \frac{9}{8\pi^2}x$ for $x \rightarrow \infty$ (see Proposition 2. in [13]), for odd integers $k \geq 7$ we have

$$(3.9) \quad |\{\text{fundamental discriminants } D \mid 0 < D < x \text{ and } 3 \nmid \beta_k(D)\}| \gg \left(\frac{9}{16\pi^2} - \epsilon\right)x.$$

Since $\Phi_{k+\frac{1}{2}}(z) \in S_{k+\frac{1}{2}}^+(\Gamma_0(4))$, the cusp form $\Phi_{k+\frac{1}{2}}(z)$ is a linear combination of Hecke eigenforms $g_\ell(z) = \sum_{n=1}^{\infty} c_\ell(n)q^n \in S_{k+\frac{1}{2}}^+(\Gamma_0(4))$ for $1 \leq \ell \leq d_k$. By Theorem 2.3 we complete the proof of Theorem.

From now on, suppose that k is even, and that $d_k > 1$ or $k = 10$. Let

$$(3.10) \quad \Psi_{k+\frac{1}{2}}(z) := \Delta(4z)R_{k-12}(z)\theta(z) = \sum_{n=1}^{\infty} \alpha_k(n)q^n \quad \text{for } k > 10$$

and

$$(3.11) \quad \begin{aligned} \Psi_{10+\frac{1}{2}}(z) &:= -(\theta(z)E_4(4z)E_6(4z) - H_{\frac{5}{2}}(z)E_4(4z)^2) \otimes \chi_3 \\ &\quad + (\theta(z)E_4(4z)E_6(4z) - H_{\frac{5}{2}}(z)E_4(4z)^2) \otimes \chi_3^2, \end{aligned}$$

where $\chi_3(n) = \left(\frac{n}{3}\right)$. We have by the sturm's bound and (3.5)

$$\begin{aligned} \sum_{n \equiv 2 \pmod{3}} \alpha_k(n)q^n &\equiv \sum_{n \equiv 2 \pmod{3}} \alpha_{12}(n)q^n \\ &\equiv 2q^8 + 2q^{17} + q^{20} + 2q^{41} + q^{44} + q^{53} + q^{56} + q^{65} + 2q^{68} + 2q^{80} + 2q^{89} + 2q^{92} \dots \\ &\equiv \sum_{n \equiv 2 \pmod{3}} H(1, 3n)q^n \pmod{3}. \end{aligned}$$

The remained part of the proof can be completed in a similar way as before, so we omit the details. $\square$

Remark 3.2. The argument given in [[10], p. 186 bottom] in the case where k is even and $k \equiv 1 \pmod{3}$ is not correct, since it would require that all the coefficients of δ_{k-4} are 3-integral which in general is not the case.

4. Conclusion

In this paper, we extend the result in [10] to the case when k is any integer, in particular when k is odd. So, we obtain that, for each integer $2k$ such that $\dim S_{2k} \geq 1$, there is a normalized Hecke eigenform F in S_{2k} satisfying $\mathcal{N}_F(x) \gg x$ ($x \rightarrow \infty$). We conclude this paper with the following remark:

Remark 4.1. (1) For each odd $k \geq 9$ and even λ such that $\lambda \geq 12$, all the coefficients of $\Phi_{k+\frac{1}{2}}$ and $\Psi_{\lambda+\frac{1}{2}}$ are 3-integral and a positive portion of these coefficients $\beta_k(n)$ and $\alpha_\lambda(n)$ is not vanishing modulo 3.

(2) Note that $k = 9$ is the minimum odd integer such that $\dim(S_{2k}) > 0$. Let

$$\Delta(z) = q \prod_{n=1}^{\infty} (1 - q^n)^{24}.$$

If we take $F(z) = \Delta(z)E_6(z)$, then $F(z)$ is the unique normalized Hecke eigenform in S_{18} . Corollary 1.3 implies that

$$\mathcal{N}_F(x) \geq \left(\frac{9}{16\pi^2} - \epsilon \right) x \quad (x \gg_{\epsilon} 0).$$

(3) The direct computation shows that if $f \in S_{k+\frac{1}{2}}^+(\Gamma_0(4))$ has integral coefficients for even k such that $d_k = 1$, then

$$f \equiv c \sum_{\substack{n \geq 1 \\ 3 \nmid n}} q^{n^2} \pmod{3}$$

for some $c \in \{-1, 1\}$.

Acknowledgement

The first author was supported by KRF-2008-331-C00005 and wish to express his gratitude to KIAS for its support through Associate membership program. The second author was partially supported by KOSEF-R01-2008-000-20446 and KRFKRF-2007-412-J02302.

REFERENCES

- [1] H. Cohen. Sums involving the values at negative integers of L -functions of quadratic characters. *Math. Ann.*, 217(3):271–285, 1975.
- [2] H. Davenport and H. Heilbronn. On the density of discriminants of cubic fields. II. *Proc. Roy. Soc. London Ser. A*, 322(1551):405–420, 1971.
- [3] D. W. Farmer and K. James. The irreducibility of some level 1 Hecke polynomials. *Math. Comp.*, 71(239):1263–1270 (electronic), 2002.
- [4] S. Friedberg and J. Hoffstein. Nonvanishing theorems for automorphic L -functions on $GL(2)$. *Ann. of Math. (2)*, 142(2):385–423, 1995.
- [5] D. Goldfeld. Conjectures on elliptic curves over quadratic fields. In *Number theory, Carbondale 1979 (Proc. Southern Illinois Conf., Southern Illinois Univ., Carbondale, Ill., 1979)*, volume 751 of *Lecture Notes in Math.*, pages 108–118. Springer, Berlin, 1979.
- [6] H. Hida and Y. Maeda. Non-abelian base change for totally real fields. *Pacific J. Math.*, (Special Issue):189–217, 1997. Olga Taussky-Todd: in memoriam.
- [7] K. James. L -series with nonzero central critical value. *J. Amer. Math. Soc.*, 11(3):635–641, 1998.
- [8] W. Kohnen. Modular forms of half-integral weight on $\Gamma_0(4)$. *Math. Ann.*, 248(3):249–266, 1980.
- [9] W. Kohnen. Fourier coefficients of modular forms of half-integral weight. *Math. Ann.*, 271(2):237–268, 1985.

- [10] W. Kohnen. On the proportion of quadratic character twists of L -functions attached to cusp forms not vanishing at the central point. *J. Reine Angew. Math.*, 508:179–187, 1999.
- [11] W. Kohnen and D. Zagier. Values of L -series of modular forms at the center of the critical strip. *Invent. Math.*, 64(2):175–198, 1981.
- [12] S. Lang, *Introduction to Modular Forms*, Grundle. d. Math. Wiss. no. 222, Springer: Berlin Heidelberg New York, 1976 Berlin, 1995.
- [13] J. Nakagawa and K. Horie. Elliptic curves with no rational points. *Proc. Amer. Math. Soc.*, 104(1):20–24, 1988.
- [14] K. Ono and C. Skinner. Fourier coefficients of half-integral weight modular forms modulo l . *Ann. of Math. (2)*, 147(2):453–470, 1998.
- [15] G. Shimura. The special values of the zeta functions associated with cusp forms. *Comm. Pure Appl. Math.*, 29(6):783–804, 1976.
- [16] J. Sturm. On the congruence of modular forms. In *Number theory (New York, 1984–1985)*, volume 1240 of *Lecture Notes in Math.*, pages 275–280. Springer, Berlin, 1987.
- [17] J. Urbanowicz. On the divisibility of generalized Bernoulli numbers. In *Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983)*, volume 55 of *Contemp. Math.*, pages 711–728. Amer. Math. Soc., Providence, RI, 1986.
- [18] J.L. Waldspurger. Sur les coefficients de Fourier des formes modulaires de poids demi-entier. *J. Math. Pures Appl. (9)*, 60(4):375–484, 1981.

SCHOOL OF LIBERAL ARTS AND SCIENCES, KOREA AEROSPACE UNIVERSITY, GOYANG, GYEONGGI 412-791, KOREA

E-mail address: choija@kau.ac.kr

DEPT OF MATHEMATICS AND PMI, POSTECH, POHANG , KOREA 790-784

E-mail address: yjc@postech.ac.kr